



IDA-TALLINNA KESKHAIGLA

AS Ida-Tallinna Keskhaigla  
Ravi 18, 10138 Tallinn  
Rg-kood 10822068  
Tel 666 1900  
E-post info@itk.ee  
www.itk.ee

## Whole-body static scintigraphy and/or SPECT with iodine

**A follow-up examination is performed no earlier than 6 months after treatment with radioactive iodine (radioiodine).**

The procedure includes a doctor's appointment, the administration of a radioiodine capsule in a diagnostic dose and, after 48 hours, a whole-body scan and three-dimensional SPECT/CT imaging of the neck and chest to assess the effectiveness of radioiodine therapy.

### **With thyrotropin; Thyrogen®**

You should follow a low-iodine diet for 3 weeks before the examination. There is no need to stop taking your thyroid medication (L-Thyroxin). For the purpose of TSH stimulation, thyrotropin (rhTSH/Thyrogen) is injected intramuscularly on two consecutive days before administering the radioiodine capsule. The follow-up examination is performed in an outpatient setting. The patient is not hospitalised.

### **With levothyroxine withdrawal**

You should stop taking your thyroid medication (L-Thyroxin) 4 weeks before the examination and follow a low-iodine diet for 3 weeks before the examination. The follow-up examination is performed in an outpatient setting. The patient is not hospitalised.